



Child Passenger Safety

Dennis R. Durbin, MD, MSCE, FAAP,^a Benjamin D. Hoffman, MD, FAAP,^b COUNCIL ON INJURY, VIOLENCE, AND POISON PREVENTION

Child passenger safety has dramatically evolved over the past decade; however, motor vehicle crashes continue to be the leading cause of death for children 4 years and older. This policy statement provides 4 evidence-based recommendations for best practices in the choice of a child restraint system to optimize safety in passenger vehicles for children from birth through adolescence: (1) rear-facing car safety seats as long as possible; (2) forward-facing car safety seats from the time they outgrow rear-facing seats for most children through at least 4 years of age; (3) belt-positioning booster seats from the time they outgrow forward-facing seats for most children through at least 8 years of age; and (4) lap and shoulder seat belts for all who have outgrown booster seats. In addition, a fifth evidence-based recommendation is for all children younger than 13 years to ride in the rear seats of vehicles. It is important to note that every transition is associated with some decrease in protection; therefore, parents should be encouraged to delay these transitions for as long as possible. These recommendations are presented in the form of an algorithm that is intended to facilitate implementation of the recommendations by pediatricians to their patients and families and should cover most situations that pediatricians will encounter in practice. The American Academy of Pediatrics urges all pediatricians to know and promote these recommendations as part of child passenger safety anticipatory guidance at every health supervision visit.

Improved vehicle crashworthiness and greater use of child restraint systems have significantly affected the safety of children in automobiles. Major shifts in child restraint use, particularly the use of booster seats among older children, have occurred in response to public education programs and enhancements to child restraint laws in nearly every state.^{1–3} In addition, there has been a substantial increase in scientific evidence on which to base recommendations for best practices in child passenger safety. Current estimates of child restraint effectiveness indicate that child safety seats reduce the risk of injury by 71% to 82%^{4, 5} and reduce the risk of death by 28% when compared with children of similar ages in seat belts.⁶ Booster seats reduce the risk of nonfatal injury among 4- to 8-year-olds by 45% compared with seat belts.⁷ Despite this

abstract

^aDepartment of Pediatrics, The Ohio State University College of Medicine and Nationwide Children's Hospital, Columbus, Ohio; and
^bDepartment of Pediatrics, Oregon Health and Science University, Portland, Oregon

Drs Durbin and Hoffman were equally responsible for conception and design of the revision, drafting, and editing of the manuscript.

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

DOI: <https://doi.org/10.1542/peds.2018-2460>

Address correspondence to Benjamin D. Hoffman, MD, FAAP. E-mail: hoffmanb@ohsu.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2018 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they do not have a financial relationship relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

To cite: Durbin DR, Hoffman BD, AAP COUNCIL ON INJURY, VIOLENCE, AND POISON PREVENTION. Child Passenger Safety. *Pediatrics*. 2018;142(5):e20182460

TABLE 1 Summary of Best Practice Recommendations

Best Practice Recommendation	Complementary Information
<p>1) Best practice recommendation: infant-only or convertible CSS used rear facing</p> <p>All infants and toddlers should ride in a rear-facing CSS as long as possible, until they reach the highest weight or height allowed by their CSS's manufacturer.</p>	<p>Rear-facing-only seats usually have a handle for carrying and can be snapped in and out of a base that is installed in the vehicle. They can only be used rear facing. Convertible CSSs can be used either forward or rear facing and typically have higher rear-facing weight and height limits than rear-facing-only seats.</p> <p>When children using rear-facing-only seats reach the highest weight for their seat, they should continue to ride rear-facing in a convertible seat for as long as possible. Most currently available convertible seats can be used rear facing to at least 40 lb.</p>
<p>2) Best practice recommendation: convertible or combination CSS used forward facing</p> <p>All children who have outgrown the rear-facing weight or height limit for their CSS should use a forward-facing CSS with a harness for as long as possible, up to the highest weight or height allowed by their CSS's manufacturer.</p>	<p>Combination CSSs are seats that can be used forward facing with a harness system and then, when the child exceeds the height or weight limit for the harness, as a booster seat with the harness removed.</p> <p>Most models of convertible and combination CSSs can accommodate children up to 65 lb and some up to 70–90 lb when used forward facing. The lowest maximum weight limit for currently available forward-facing car safety seats is 40 lb.</p>
<p>3) Best practice recommendation: belt-positioning booster seat</p> <p>All children whose weight or height is above the forward-facing limit for their CSS should use a belt-positioning booster seat until the vehicle lap and shoulder seat belt fits properly, typically when they have reached 4 ft 9 inches in height and are between 8 and 12 y of age.</p>	<p>A few vehicle models offer integrated forward-facing seats with a harness system. The vehicle owner's manual provides instructions for use of integrated seats when they are present. A crash-tested travel vest may be considered for children with special needs or in situations where a traditional CSS cannot be installed correctly.</p> <p>There is a safety advantage for young children to remain in car safety seats with a harness for as long as possible before transitioning to booster seats.</p> <p>Booster seats function by positioning the child so that both the lap and shoulder portions of the vehicle seat belt fit properly: the lap portion of the belt should fit low across the hips and pelvis, and the shoulder portion should fit across the middle of the shoulder and chest. They come in both high-back (a seat back that extends up beyond the child's head) and backless models. A few vehicle models offer integrated booster seats.</p>
<p>4) Best practice recommendation: Lap and shoulder vehicle seat belt</p> <p>When children are old enough and large enough to use the vehicle seat belt alone, they should always use lap and shoulder seat belts for optimal protection</p>	<p>The lap portion of the belt should fit low across the hips and pelvis, and the shoulder portion should fit across the middle of the shoulder and chest when the child sits with his back against the vehicle seat back. If they don't, the child is likely too small to use the vehicle seat belt alone and should continue to use a belt-positioning booster seat.</p>
<p>5) Best practice recommendation: all children <13 years of age should be restrained in the rear seats of vehicles for optimal protection</p> <p>All children <13 y should be restrained in the rear seats of vehicles for optimal protection</p>	<p>CSSs should be installed tightly either with the vehicle seat belt or with the LATCH system, if available. LATCH is a system of attaching a CSS to the vehicle that does not use the seat belt. It was designed to ease installation of the CSS. Whether parents use LATCH or the seat belt, they should always ensure a tight installation of the CSS into the vehicle.</p>

LATCH, Lower Anchors and Tethers for Children.

progress, each year, nearly 1000 children younger than 16 years die in motor vehicle crashes in the United States.⁸

The American Academy of Pediatrics (AAP) strongly supports optimal safety for children and adolescents of all ages during all forms of travel. This policy statement provides 5 evidence-based recommendations for best practices to optimize safety in passenger vehicles for all children, from birth through adolescence (summary of recommendations in Table 1):

1. All infants and toddlers should ride in a rear-facing car safety seat

(CSS) as long as possible, until they reach the highest weight or height allowed by their CSS's manufacturer. Most convertible seats have limits that will permit children to ride rear-facing for 2 years or more.

2. All children who have outgrown the rear-facing weight or height limit for their CSS should use a forward-facing CSS with a harness for as long as possible, up to the highest weight or height allowed by their CSS's manufacturer.

3. All children whose weight or height is above the forward-facing limit for their CSS should use a

belt-positioning booster seat until the vehicle lap and shoulder seat belt fits properly, typically when they have reached 4 ft 9 inches in height and are between 8 and 12 years of age.

4. When children are old enough and large enough to use the vehicle seat belt alone, they should always use lap and shoulder seat belts for optimal protection.

5. All children younger than 13 years should be restrained in the rear seats of vehicles for optimal protection.




FIGURE 1

Algorithm to guide implementation of best practice recommendations for optimal child passenger safety. (See Table 1 for summary of recommendations and Table 2 for definitions and explanations.)

TABLE 2 Explanations of Decision Points and Additional Resources

Does the child have significant health needs?	Children with certain temporary or permanent physical and behavioral conditions such as altered muscle tone, decreased neurologic control, skeletal abnormalities, or airway compromise that may preclude the use of regular CSSs may require specialized restraint systems.
Consult complementary AAP Policy and other resources for best practice recommendations	The AAP has a Policy Statement providing specific guidance on best practice recommendations for children with special health care needs (http://aappolicy.aappublications.org/cgi/content/full/pediatrics%3B104/4/988). To locate a child passenger safety technician in your area with special training in special health needs, go to http://cert.safekids.org .
<4 y	Infants and toddlers have relatively large heads and several structural features of their neck and spine that place them at particularly high risk of head and spine injuries in motor vehicle crashes. Rear-facing CSSs provide optimal support to the head and spine in the event of a crash. Children who are small for their age may need to be evaluated like younger children. Consult a child passenger safety technician with enhanced training in special needs or other resources for assistance.
Has the child outgrown weight or height limit for seat?	The AAP annually updates information on child restraint systems currently available in the United States (www.healthychildren.org/carseatguide). The weight thresholds provided in the algorithm are considered minimum standards. More recent products have higher weight limits and should be used when possible. In general, children should remain in a child restraint system until they outgrow the weight or height limits for its intended use.
Is weight or height less than rear-facing limit for convertible CSS?	
Is weight or height less than forward-facing limit for convertible combination CSS?	
4–8 y	Most children 4–8 years of age are not large enough to fit properly in the vehicle seat belt and will require a CSS or booster seat for optimal restraint. A belt-positioning booster seat positions a child so that the lap and shoulder portions of the seat belt fit properly: the lap portion low across the hips and pelvis and the shoulder portion across the middle of the shoulder and chest.
8 y	Most children under 4 ft 9 inches in height will not fit properly in vehicle lap and shoulder seat belts.
Does child fit properly in the vehicle seat belt, usually around 4 ft 9 inches in height?	These 3 questions are an evaluation to determine whether a child is ready to be restrained by the vehicle seat belt without a booster seat. If the answer is “no” to any of these questions, the child should use a booster seat: 1. Is the child tall enough to sit against the vehicle seat back with her knees bent at the edge of the vehicle seat without slouching and stay in this position comfortably throughout the trip? 2. Does the shoulder belt lie across the middle of the chest and shoulder, not against the neck or face? 3. Is the lap belt low across the hips and pelvis?

Of note, the recommendation that all children be restrained in a rear-facing-only or convertible CSS used rear facing as long as possible represents a significant change from previous AAP policy and is based on data from the United States⁹ as well as extensive experience in Sweden.^{10,11} It is important to note that nearly all currently available CSSs have weight limits for rear-facing use that can accommodate children 35 to 40 lb.¹²

Certain considerations contained in this policy statement are relevant to commercial airline travel as well and are noted in the accompanying technical report. Additional AAP policy statements provide specific recommendations to optimize safety

for preterm and low birth weight infants,¹³ children in school buses,¹⁴ and children using other forms of travel and recreational vehicles.^{15–17} In addition, complementary AAP policy statements provide recommendations for teen drivers¹⁸ and for the safe transport of newborn infants¹⁹ and children with special health care needs.^{20,21}

Pediatricians play a critical role in promoting child passenger safety. To facilitate their widespread implementation in practice, evidence-based recommendations for optimal protection of children of all ages in passenger vehicles are presented in the form of an algorithm (Fig 1) with an accompanying table of explanations and definitions (Table

2). A summary of the evidence in support of these recommendations is provided in the accompanying technical report.²² Because pediatricians are a trusted source of information to parents, pediatricians need to maintain a basic level of knowledge of these best practice recommendations and promote and document them at every health supervision visit. Prevention of motor vehicle crash injury is unique in health supervision topics, as it is the only topic recommended at every health supervision visit by *Bright Futures*.²³ Pediatricians can also use this information to promote child passenger safety public education, legislation, and regulation at local, state, and national levels through

a variety of advocacy activities, including aligning their state's child passenger safety law with the best practice recommendations promoted in this policy statement.

Pediatricians are urged to keep abreast of the evolving and multifaceted guidance and resources on motor vehicle safety for children. In particular, many communities have child passenger safety technicians who have completed a standardized National Highway Traffic Safety Administration course and who can provide hands-on advice and guidance to families. In most communities, child passenger safety technicians work at formal inspection stations; a list of these is available at <https://www.nhtsa.gov/equipment/car-seats-and-booster-seats#install-inspection>. If your community does not have an inspection station, you can find a child passenger safety technician in your area via the National Child Passenger Safety Certification Web site at <http://cert.safekids.org> or via the National Highway Traffic Safety Administration Child Safety Seat Inspection Station Locator at <http://www.nhtsa.dot.gov/cps/cpsfitting/index.cfm>. Car Seat Check Up Events are updated at https://www.safekids.org/events/field_type/check-event. In addition, additional resources for pediatricians and families can be found at www.aap.org or www.healthychildren.org.

LEAD AUTHORS

Dennis R. Durbin, MD, MSCE, FAAP

Benjamin D. Hoffman, MD, FAAP

COUNCIL ON INJURY, VIOLENCE AND POISON PREVENTION, 2018–2019

Benjamin D. Hoffman, MD, FAAP, Chairperson

Phyllis F. Agran, MD, MPH, FAAP

Sarah A. Denny, MD, FAAP

Michael Hirsh, MD, FAAP

Brian Johnston, MD, MPH, FAAP

Lois K. Lee, MD, MPH, FAAP

Kathy Monroe, MD, FAAP

Judy Schaechter, MD, MBA, FAAP

Milton Tenenbein, MD, FAAP

Mark R. Zonfrillo, MD, MSCE, FAAP

Kyran Quinlan, MD, MPH, FAAP, ex-officio

CONTRIBUTOR

Stuart Weinberg, MD, FAAP, Partnership for Policy Implementation

LIAISONS

Lynne Janecek Haverkos, MD, MPH, FAAP – *National Institute of Child Health and Human Development*

Jonathan D. Midgett, PhD – *Consumer Product Safety Commission*

Alexander W. (Sandy) Sinclair – *National Highway Traffic Safety Administration*

Richard Stanwick, MD, FAAP – *Canadian Paediatric Society*

STAFF

Bonnie Koziol

ABBREVIATIONS

AAP: American Academy of Pediatrics

CSS: car safety seat

REFERENCES

1. Durbin DR, Kallan MJ, Winston FK. Trends in booster seat use among young children in crashes. *Pediatrics*. 2001;108(6). Available at: www.pediatrics.org/cgi/content/full/108/6/e109
2. Winston FK, Chen IG, Elliott MR, Arbogast KB, Durbin DR. Recent trends in child restraint practices in the United States. *Pediatrics*. 2004;113(5). Available at: www.pediatrics.org/cgi/content/full/113/5/e458
3. Insurance Institute for Highway Safety. Child Restraint/Belt Use Laws. Arlington, VA: Insurance Institute for Highway Safety, Highway Loss Data Institute. Available at: <http://www.iihs.org/iihs/topics/laws/safetybeltuse>. Accessed August 17, 2010
4. Arbogast KB, Durbin DR, Cornejo RA, Kallan MJ, Winston FK. An evaluation of the effectiveness of forward facing child restraint systems. *Accid Anal Prev*. 2004;36(4):585–589
5. Zaloshnja E, Miller TR, Hendrie D. Effectiveness of child safety seats vs

safety belts for children aged 2 to 3 years. *Arch Pediatr Adolesc Med*. 2007;161(1):65–68

6. Elliott MR, Kallan MJ, Durbin DR, Winston FK. Effectiveness of child safety seats vs seat belts in reducing risk for death in children in passenger vehicle crashes. *Arch Pediatr Adolesc Med*. 2006;160(6):617–621
7. Arbogast KB, Jermakian JS, Kallan MJ, Durbin DR. Effectiveness of belt positioning booster seats: an updated assessment. *Pediatrics*. 2009;124(5):1281–1286
8. National Highway Traffic Safety Administration. Fatality analysis reporting system (FARS) encyclopedia. Available at: <http://www.fars.nhtsa.dot.gov/>. Accessed August 21, 2018
9. McMurry TL, Arbogast KB, Sherwood CP, et al. Rear-facing versus forward-facing child restraints: an updated assessment. *Inj Prev*. 2018;24(1):55–59
10. Isaksson-Hellman I, Jakobsson L, Gustafsson C, Norin HA. Trends and effects of child restraint systems based on Volvo's Swedish accident database. In: *Proceedings of Child Occupant Protection 2nd Symposium*; November 12, 1997; Warrendale, PA
11. Jakobsson L, Isaksson-Hellman I, Lundell B. Safety for the growing child: experiences from Swedish accident data. In: *Proceedings from the 19th International Technical Conference on the Enhanced Safety of Vehicles*; June 6–9, 2005; Washington, DC. Abstr 05-0330
12. American Academy of Pediatrics. *Car Safety Seats: Product Listing for 2018*. Elk Grove Village, IL: Healthy Children, American Academy of Pediatrics; 2018. Available at: www.healthychildren.org/carseatlist. Accessed June 8, 2018
13. Bull MJ, Engle WA; Committee on Injury, Violence, and Poison Prevention and Committee on Fetus and Newborn; American Academy of Pediatrics. Safe transportation of preterm and low birth weight infants at hospital discharge. *Pediatrics*. 2009;123(5):1424–1429
14. Agran PF; American Academy of Pediatrics Committee on Injury, Violence, and Poison Prevention; American Academy of Pediatrics

- Council on School Health. School transportation safety. *Pediatrics*. 2007;120(1):213–220
15. American Academy of Pediatrics, Committee on Injury and Poison. All-terrain vehicle injury prevention: two-, three-, and four-wheeled unlicensed motor vehicles. *Pediatrics*. 2000;105(6):1352–1354
 16. American Academy of Pediatrics, Committee on Injury and Poison Prevention. Bicycle helmets. *Pediatrics*. 2001;108(4):1030–1032
 17. American Academy of Pediatrics, Committee on Injury and Poison Prevention. Personal watercraft use by children and adolescents. *Pediatrics*. 2000;105(2):452–453
 18. Weiss JC; Committee on Injury, Violence, and Poison Prevention, American Academy of Pediatrics; Committee on Adolescence, American Academy of Pediatrics. The teen driver. *Pediatrics*. 2006;118(6):2570–2581
 19. Bull M, Agran P, Laraque D, et al; American Academy of Pediatrics, Committee on Injury and Poison Prevention. Safe transportation of newborns at hospital discharge. *Pediatrics*. 1999;104(4, pt 1):986–987
 20. O’Neil J, Hoffman BD; Council on Injury, Violence, and Poison Prevention. School bus transportation of children with special health care needs [published correction appears in *Pediatrics*. 2018;142(1):e20181221]. *Pediatrics*. 2018;141(5):e20180513
 21. Bull M, Agran P, Laraque D, et al; American Academy of Pediatrics, Committee on Injury and Poison Prevention. Transporting children with special health care needs. *Pediatrics*. 1999;104(4, pt 1):988–992
 22. Durbin DR, Hoffman B; American Academy of Pediatrics, Committee on Injury, Violence, and Poison Prevention. Child passenger safety. *Pediatrics*. 2018;142(5):e20182461
 23. American Academy of Pediatrics. Bright futures: health care professionals tools and resources web site. Available at: <http://brightfutures.aap.org/materials-and-tools/guidelines-and-pocket-guide/Pages/default.aspx>. Accessed August 17, 2010

Child Passenger Safety

Dennis R. Durbin, Benjamin D. Hoffman and COUNCIL ON INJURY, VIOLENCE,
AND POISON PREVENTION

Pediatrics originally published online August 30, 2018;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/early/2018/08/28/peds.2018-2460
References	This article cites 17 articles, 13 of which you can access for free at: http://pediatrics.aappublications.org/content/early/2018/08/28/peds.2018-2460#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Current Policy http://www.aappublications.org/cgi/collection/current_policy Council on Injury, Violence, and Poison Prevention http://www.aappublications.org/cgi/collection/committee_on_injury_violence_and_poison_prevention Injury, Violence & Poison Prevention http://www.aappublications.org/cgi/collection/injury_violence_-_poison_prevention_sub Carseat Safety http://www.aappublications.org/cgi/collection/carseat_safety_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Child Passenger Safety

Dennis R. Durbin, Benjamin D. Hoffman and COUNCIL ON INJURY, VIOLENCE,
AND POISON PREVENTION

Pediatrics originally published online August 30, 2018;

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2018/08/28/peds.2018-2460>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2018 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

